

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Macromolecular and Micronutrient Profiles of Sprouted Chickpeas to Be Used for Integrating Cereal-Based Food

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1731154> since 2021-09-28T18:13:36Z

Published version:

DOI:10.1094/CCHEM-04-16-0108-FI

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Macromolecular and micronutrient profiles of sprouted chickpeas to be used for integrating cereal-based food

Journal:	<i>Cereal Chemistry</i>
Manuscript ID	Draft
Manuscript Type:	Focus Issue
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Marengo, Mauro; Università degli Studi di Milano, DeFENS, Department of Food, Nutritional and Environmental Sciences Carpen, Aristodemo; Università degli Studi di Milano Bonomi, Francesco; University of Milan, Casiraghi, Maria Cristina; University of Milan, DeFENS - Department of Food Environmental and Nutritional Sciences Meroni, Erika; University of Milan, Department of Food, Environmental and Nutritional Sciences Quaglia, Lucio; Molino Quaglia Iametti, Stefania; University of Milan, DISMA; Pagani, M. Ambrogina; Università degli Studi di Milano, DeFENS, Department of Food, Nutritional and Environmental Sciences Marti, Alessandra; Università degli Studi di Milano, DeFENS, Department of Food, Nutritional and Environmental Sciences</p>
Area of Expertise:	Dough, Oilseeds and Legumes, Protein

**Macromolecular and micronutrient profiles of sprouted chickpeas to be used for
integrating cereal-based food**

Mauro Marengo¹, Aristodemo Carpen¹, Francesco Bonomi¹, Maria Cristina Casiraghi¹, Erika Meroni¹,
Lucio Quaglia², Stefania Iametti¹, Maria Ambrogina Pagani^{§,1}, Alessandra Marti¹

¹Dipartimento di Scienze per gli Alimenti, la Nutrizione e l’Ambiente (DeFENS), Università degli Studi di
Milano, Italy; ²Molino Quaglia, Vighizzolo d’Este, Italy

[§]Corresponding author
ambrogina.pagani@unimi.it

DeFENS
Via G. Celoria, 2
20133 Milan, Italy
Phone +3902-5031-6658

ABSTRACT

Pulse flour may be used to improve nutritional traits of gluten and gluten-free formulations in traditional food such as bread or pasta. However due some intrinsic nutritional, textural, and sensory properties, the use of pulses as ingredients for production of enriched food remains limited. In this study, we investigated the modification in macromolecules and micronutrients in industrial-scale flour from partially sprouted chickpeas to define its possible use as an ingredient in cereal-based foods. Controlled sprouting resulted in significant decrease of antinutritional compounds and in an increase of free minerals and vitamins. Sprouting also affected the overall structural organization of proteins and their thiol/disulfide balance, and promoted release of peptides. All of these had a positive effect on dough mixing properties, in particular for dough development. Formulations with enrichment in sprouted chickpeas flour (wheat:chickpea ratio = 100:20) were tested also as for their dough leavening properties that improved with respect to flour from non-sprouted chickpeas. All together, taking into account the modifications induced by partial sprouting at industrial scale, we can conclude that sprouted chickpea flour represents an interesting ingredient for production of enriched cereal based food with better nutritional and rheological characteristics.

Keywords: chickpeas, sprouting, thiol accessibility, pasting properties, dough rheology

1
2
3 34 Pulses are a particularly rich source of vegetable proteins, as well as of dietary fibers, vitamins and minerals
4
5 35 (such as folate and iron), but their nutritional value is limited by the presence of antinutrients that decrease
6
7 36 digestibility and micronutrient bioavailability (Oomah et al 2011). Thus, despite Western consumers are
8
9 37 increasingly interested in natural and healthy food products – without turning down the hedonistic aspect
10
11 38 of food – the use of pulses are underexploited at an industrial scale, and they are seldom used as
12
13 39 ingredients in processed foods such as bakery products, pasta and snacks (Abu-Ghannam and Gowen,
14
15 40 2011). Indeed, in addition to nutritional issues, consumers often identify off-flavors in products containing
16
17 41 high levels (>10%) of pulse flours.

18
19
20 42 Sprouting (or germination) is a natural process that decreases the antinutrient content in pulses
21
22 43 while substantially increases the micronutrient bioavailability and improving the sensory properties. The
23
24 44 nutritional benefits of germination have been documented extensively in the scientific literature (Savelkoul
25
26 45 et al 1992; Chitra et al 1996; Vidal-Valverde et al 2002; Ghavidel and Prakash 2007; Martín-Cabrejas et al
27
28 46 2008; Khandelwal et al 2010; Gujral et al 2011). During germination, natural enzymes that degrade
29
30 47 antinutrients such as phytic acid and trypsin inhibitors are expressed, increasing the vitamin content and
31
32 48 the bioavailability of some minerals in the products. Enzymes are also able to degrade ROF (raffinose family
33
34 49 of oligosaccharides) into shorter carbohydrates, decreasing the digestive discomfort caused by the
35
36 50 presence of ROF and developing a sweet flavor note in germinated pulses.

37
38
39 51 In many countries, the germination process traditionally has been performed at a household level.
40
41 52 The basic germination process consists of steeping pulses in water until they reach the moisture content
42
43 53 needed to initiate the seedling, after which the steeping water is drained and the pulses are allowed to
44
45 54 germinate under cool, humid conditions. After the germination phase, the grains need to be dried since
46
47 55 they usually have a moisture content of 30-50%. Processing conditions must be constantly and precisely
48
49 56 monitored and controlled to standardize the properties of the final product and assure good quality is
50
51 57 maintained (Bellaio et al 2014).

52
53
54 58 Some studies had shown that partial germination of several types of pulses decreased the
55
56 59 antinutrient content and improved the amount of micronutrients (Ghavidel et al 2007; Bellaio et al 2012).
57
58 60 Moreover, the content of fructose greatly increased, thus improving the ingredient taste with a desirable
59
60

sweet note (Bellaio et al 2012). In addition to nutritional aspects, germination positively affected the dehulling process for brown chickpeas, mung peas, and pigeon peas (Bellaio et al 2011). Moreover, the process can influence the cooking properties of pulses, including cooking time and dispersed solids (Zamproga et al 2011). Finally, developing processed foods such as pasta, white layer cakes, and extruded snacks with partially germinated yellow pea reportedly resulted in end products with acceptable characteristics (Han and Buchko 2014; Frohlinch et al 2014).

The impact of sprouting – carried out at an industrial scale – on macromolecular and micronutrient profiles of chickpeas was assessed in the present study. Modification of protein overall association as well variation of specific proteins contents and starch pasting properties were investigated before and after sprouting. In addition, rheological properties of chickpea flour enriched dough were studied in view of providing a nutritious and high protein staple food for low-income end users, such as those in developing areas, and for users wishing to improve the nutritional quality of their diet.

MATERIALS AND METHODS

Materials. Raw and sprouted chickpea (Fig. S1) were provided by Molino Quaglia (Vighizzolo d'Este, Italy). Chickpeas were sprouted in an industrial sprouting plant (Bühler AG, Uzwil, Switzerland) for 3 days at 18-24 °C. Sprouted peas were dried at 50 °C, and stored at room temperature. When required, chickpeas samples (sprouted or non-sprouted) were ground into flour (0.5 mm particle size) in a pin mill (Bühler AG, Uzwil, Switzerland). A commercial common wheat flour (proteins: 13%; alveographic W: 355 *10⁻⁴J; alveographic P/L: 0.76) was used to prepare the chickpea-enriched dough for rheology studies (wheat:chickpea ratio = 100:20).

Chemical Composition. Moisture, starch, protein, lipid and ash content was assessed by AACC standard methods (44-15.02, 76-13.01, 46-12.01, 30-10.01, and 08-01.01, respectively; AACC 2001). Sugars were assessed by HPLC by Anion Exchange Chromatography with Pulsed Amperometric Detection (HPAEC-PAD) (Zygmunt et al 1982). Total, soluble and insoluble dietary fiber content was by the enzymatic–gravimetric

procedure (AOAC Method 991.43). The analysis of phytic acid was performed by HPLC with method that uses a spectrophotometric detection as described by Oberleas & Harland (2007).

Protein solubility, soluble peptides, and thiol accessibility. The solubility of proteins in chickpea flour samples (10 g) was determined by suspending the flour samples (10 g in 10 L⁻¹) in 0.05 mol L⁻¹mM sodium phosphate, 0.1 mol L⁻¹ NaCl buffer, pH 7.0, containing 8 mol L⁻¹ urea or 8 mol L⁻¹ urea and 0.01 mol L⁻¹ dithiothreitol (DTT) where indicated (Bonomi et al 2012). After 1 h stirring at 25°C, the suspensions were centrifuged (~2,500 x g, 30 min, 25°C), and the protein concentration in the supernatant was determined by a dye-binding method (Bradford 1976).

The amount of soluble peptides was monitored by measuring the absorbance at 280 nm in the supernatant obtained from treatment with 10% trichloroacetic acid (TCA). Typically, flour (10 g) was suspended in 10 mL of 0.05 mol L⁻¹sodium phosphate, 0.1 mol L⁻¹NaCl buffer, pH 7.0. After stirring for 30 min at room temperature, the supernatant was collected by centrifugation at 10000 x g for 30 min, and an appropriate aliquot was treated with an equal volume of 20% TCA. The supernatant of TCA precipitation was collected by centrifugation at 10000 x g for 30 min, and read at 280 nm. Results are given as arbitrary units (absorbance at 280 nm of the TCA supernatant, corrected for dilution when necessary).

Accessible thiols (expressed as μmol thiols (g flour)⁻¹) were determined by suspending chickpea flour in 0.05 mol L⁻¹mM sodium phosphate, 0.1 mol L⁻¹NaCl buffer, pH 7.0, containing 0.2 mmol L⁻¹ 5,5'-dithiobis-(2-nitrobenzoate) (DTNB), in the presence/absence of 8 mol L⁻¹ urea as described by Bonomi et al. 2012. After 1 h stirring at 25°C and centrifugation (~2,500 x g, 30 min, 25°C), absorbance of the supernatant was read at 412 nm.

SDS-PAGE. Proteins solubilized from chickpea flour as described above were diluted to 1/10 of their volume with SDS-PAGE denaturing buffer (0.125 mol L⁻¹Tris-HCl, pH 6.8, 500 mL glycerol L⁻¹, 17 g L⁻¹ SDS, 0.1 g L⁻¹ Bromophenol Blue), containing 10 mL L⁻¹ of 2-mercaptoethanol when indicated, and the mixtures were heated at 100°C for 10 min. Sample volumes were adjusted to load 0.01 mg of protein per lane. SDS-PAGE

was carried out in a MiniProtein apparatus (BioRad, Richmond, VA, USA), and gels were stained with Coomassie Blue.

Trypsin inhibitor levels. Trypsin inhibitor content was assessed by using the official method IS/ISO 14902 (2001) as described in Scarafoni et al 2008, using 0.2 mmol L⁻¹ solutions of the chromogenic substrate N-benzoyl-p-nitroanilide (BAPA) in 150 mM Tris/HCl, pH 8.0, and trypsin (E.C.3.4.21.4 TPCK-treated, T1426, SIGMA Aldrich, Milan, Italy). Trypsin activity was measured in the presence/absence of the buffer-soluble protein fraction prepared from chickpea flour as described above. One trypsin inhibitor unit (TIU) was defined as the amount of inhibitor that decreases trypsin activity by 50%.

Enzyme Activity Assay. Proteolytic activity was determined by the method of Arnon (1970) using azocasein (Sigma Chemical Co., St Louis, MO, USA) as the substrate. Alpha-amylase activity was determined at least in duplicate according to AACC standard method n. 303, by using the Megazyme Amylase Assay Procedure (Megazyme International Ireland Ltd., Wicklow, Ireland).

Micronutrients determination. For metal quantitative analysis, samples were analyzed by ICP-MS (Bruker AURORA M90 ICP-MS, Milan, Italy). Total metal content was measured on aliquots of a flour suspension (10 g of flour suspended in 10 mL of 100 mmol L⁻¹ Tris-HCl buffer, pH 7.5) after removing solids by centrifugation at 10000 x g for 30 min. Unbound metals were assessed on the ultrafiltrate obtained by passing the clarified extract through a Centricon ultrafiltration device (5kDa nominal cutoff, Merck Millipore, Vimodrone, Italy).

A RP-HPLC method was used for the determination of water-soluble vitamins (B1, B2, B6, PP) according to (Aslam et al 2008), using a Symmetry® C18 3.5 column (5 µm, 4.6 x 250 mm (Waters, Vimodrone, Italy) and isocratic elution (water/methanol, 50/50 v/v, containing 0.2 % v/v PICB8) at a flow rate of 0.700 mL/min. The eluate was monitored at 440 and 256 nm, and vitamin content was calibrated with reference to appropriate standard curves. Vitamin determination was carried out on the supernatant obtained after centrifugation (10000 x g for 30 min) of a suspension of 10 g flour in 10 mL of 0.5 mol L⁻¹ sodium acetate pH

1
2
3 141 5.0. Unbound vitamins were assessed on the ultrafiltrate obtained by passing the clarified extract through a
4
5 142 Centricon ultrafiltration device (5kDa nominal cutoff, Merck Millipore, Vimodrone, Italy).
6
7 143
8
9 144 **Pasting Properties.** Pasting properties were measured in duplicate using a Micro-Visco-Amylograph (MVAG,
10
11 145 Brabender GmbH & Co. KG, Duisburg, Germany), as reported by Marti et al 2013, with little modification.
12
13 146 An aliquot of sample (15 g) was dispersed in 100 mL of distilled water (scaling both flour and water weight
14
15
16 147 on a 14% flour moisture basis) and stirred at 250 rpm. The following temperature profile was applied:
17
18 148 heating from 30 °C to 95 °C at a rate of 3 °C/min, holding at 95°C for 15 min, cooling from 95°C to 50°C at a
19
20 149 cooling rate of 3°C/min, and holding at 50°C for 1 min.
21
22 150
23
24 151 **Mixing Properties.** Mixing properties of chickpea-enriched doughs (wheat:chickpea ratio = 100:20) were
25
26 152 measured in duplicate using a Farinograph-E (Brabender GmbH & Co. KG, Duisburg, Germany)equipped
27
28 153 with a 50g mixing bowl. AACCI Approved Method 54-21.01 (AACC 2001) was used for the identification of
29
30
31 154 optimal water absorption for flours.
32
33 155
34
35 156 **Leavening Properties.** Leavening properties of chickpea-enriched doughs (wheat:chickpea ratio = 100:20)
36
37 157 were assessed in duplicate with a Rheofermentometer ®(Chopin, Tripette & Renaud, Villeneuve La Garenne
38
39 158 Cedex, France). Dough samples were prepared in an automatic spiral mixer (Bomann, Clatronic s.r.l.,
40
41 159 Piadena, Italy) with 1% NaCl and 1.5% bakers' yeast. Mixing time and amount of water were those
42
43
44 160 determined by the Farinograph test. The rheofermentographic test was performed on 315 g portion of the
45
46 161 dough and carried out at 30 °C for 3 h.
47
48 162
49

50
51 163 **RESULTS AND DISCUSSION**
52
53 164

54
55 165 **Sprouting influences the chemical composition.** The processing conditions used did not promote
56
57 166 important modification in terms of starch, protein (as detected as total nitrogen), lipid and fiber content
58
59 167 (Table I). In these chickpea samples the content of fiber did not change despite of many studies have shown
60

that the germination process has a significant impact on dietary fiber fractions (Chang et al 2006; Mahadevamma and Tharanathan 2004; Martin-Cabrejas et al 2003, 2008; Duenas et al 2016), but those changes are dependent on the investigated legume and germination conditions. Although the total amount of sugars did not change, sprouting promoted an important increase in sucrose and a decrease in both raffinose and glucose.

The amount of some free micronutrient as well some antinutritional compounds are positively modified by sprouting. The amount of some free micronutrient as well of some antinutritional compounds are positively modified by sprouting. Unbound metal concentrations are reported in Table II. Sprouting has an effect on the availability of some mineral micronutrients that are released from macro and micromolecules (such as phosphates and or phytate) during the process. This improvement is particularly noticeable for relevant micronutrients such as iron, copper and calcium. In this frame it is important to note that the total content of these metals did not significantly change upon sprouting (not shown), suggesting that sprouting promotes a change in speciation of these chemical species. The content of acid phytic decreases $\approx 10\%$ upon sprouting, as observed in previous studies (Chitra et al 1996). Phytates have a marked chelating ability - in particular for calcium - and interact with proteins decreasing their solubility and availability. Phytic acid has also been linked to the inhibition of digestive enzymes such as protease, alpha amylases and trypsin.

The content of unbound soluble vitamins such as B₂, B₆ and PP is not modified, while the amount of vitamin B₁ increases twice with respect to the non-sprouted sample. Sprouting lowers the levels of Serin-Protease Inhibitors (SERPINs) that are antinutrient factors broadly present in legumes. The data presented in Fig. 1 show that sprouted flour had a TIU of 42.6 ± 3 (that is, 43% less than the non-sprouted flour (24.1 ± 6 TIU)), confirming previous findings reviewed by Jain et al (2009).

Sprouting results in a modification of protein overall association as well in the content of specific proteins. Protein solubility studies in different media represents an useful approach for gathering information on the nature of the inter-protein interactions in legumes (Manful et al 2014) and in cereal-based materials (Iametti et al 2006; Bonomi et al 2012; Cabrera-Chávez et al 2012; Barbiroli et al 2013).

1
2
3 195 Sprouting resulted in a significant decrease of all the classes of proteins solubilized in the various media.
4
5 196 Proteins soluble in buffered saline were the most degraded (33.3% decrease, from 141 ± 5.4 to 94 ± 3.8 mg (g
6
7 197 flour)⁻¹), along with proteins that could only be solubilized upon treatment with dissociating agents (27.5%
8
9 198 decrease, from 218 ± 8.2 to 159 ± 7.4 mg (g flour)⁻¹), whereas changes affecting proteins in aggregates
10
11 199 stabilized only by hydrophobic bonds were much lower (14.9%, from 152 ± 7.1 to 129 ± 5.4 mg (g flour)⁻¹).
12

13 200 The electrophoretic profile of proteins and large peptides extracted in media of various dissociating
14
15 201 ability and analyzed in the absence and in the presence of disulfide-reducing agents are shown in the two
16
17 202 panels of Fig. 2, and indicate that the most evident germination-related changes involved proteins of high
18
19 203 molecular mass, that are present in the original flour as hydrophobic aggregates stabilized by disulfide
20
21 204 bonds. Recently, Ghumman et al (2016) shoed that pulses germination resulted in the breakdown of high
22
23 205 molecular weight proteins to low molecular weight proteins. A comparison between the solubility data
24
25 206 presented in the previous paragraph and the tracings in Fig. 2 indicates that small peptides (escaping
26
27 207 detection by the dye-binding method used here, and not evident in the SDS-PAGE gels) were possibly the
28
29 208 most abundant products of sprouting-induced protein breakdown.
30
31

32 209 This interpretation is supported by measurements of the TCA-soluble material in extracts obtained
33
34 210 from chickpea flours with buffered saline. Small peptides are expected to be soluble in this medium, and to
35
36 211 remain soluble even in the presence of 10% TCA. We found a $\approx 60\%$ increase in the amount of TCA-soluble
37
38 212 peptides in the flour obtained from sprouted chickpeas. The release of peptides during sprouting
39
40 213 represents a positive effect, as it may improve protein digestibility, facilitate further degradation of
41
42 214 proteins and production of bioactive species, and provide substrates for flavor formation upon further
43
44 215 processing and baking.
45
46

47 216 The formation of a proper protein network in dough has been shown to depend on the presence
48
49 217 and/or formation of intra- and intermolecular S-S bonds among involved proteins. Thus, sulfhydryls (-SH)
50
51 218 and disulfides (-S-S-) have a fundamental role in defining the technological properties of flours.
52
53 219 Quantification of the amount and accessibility of protein -SH groups has been applied to understand the
54
55 220 nature and to evaluate the extent of process-related modifications in many cereal-based foods (Bonomi et
56
57 221 al 2012; Cabrera-Chávez et al 2012; Barbiroli et al 2013). The arrangement of thiols in the different chickpea
58
59
60

samples was studied by assessing the amount of –SH groups accessible to the thiol reagent, DTNB, in the presence/absence of denaturants. Sprouting had no effect on the content of readily-accessible thiols (i.e., reacting with DTNB in the absence of denaturants) that remained constant at 120 ± 8 nmol (g flour)⁻¹ in both samples. Treatment with urea increased the content in accessible thiols to 150 ± 11 nmol (g flour)⁻¹ in the flour from non-sprouted chickpeas, whereas the content in urea-accessible thiols in flour from sprouted chickpeas was in excess of 350 nmol (g flour)⁻¹. Thus, the sprouting treatment may imply a loosening of the compact structure of legume storage proteins, making a higher number of protein thiols accessible upon protein unfolding. This new structural organization could facilitate the incorporation of these proteins into the protein network relevant in baked foods, and also improve the network-forming ability in dough containing mixtures of flour from different sources.

Sprouting also affected the endogenous levels of hydrolytic activities. The most notable increase involved alpha-amylase, that in the sprouted material was 0.125 ± 0.02 U (g flour)⁻¹, that is more than twice the activity in raw chickpea flour. The endogenous proteolytic activity also reached 7.1 ± 0.04 U/(g flour)⁻¹ in the flour from sprouted chickpea, 170% higher than in the non-sprouted sample. The presence of these endogenous enzymes may represent a form of added value, since sprouted materials can be used as partial or even total replacement for commonly used exogenous enzymes. This might allow to produce "clean label" products.

Sprouting does not compromise starch pasting properties. Sprouting affects pasting properties of chickpea flour (Fig. 3), decreasing the maximum viscosity reached during the heating step (from 416.5 to 339.5 BU for raw and sprouted chickpea flour, respectively). This is likely due to the alpha-amylase activity, which increases in sprouted chickpea. The action of α -amylase results in a rapid decrease in the size of the starchy molecules and, consequently, in the viscosity of the suspension (Collado and Corke 1999). A quite similar behavior was also observed on sprouted grains (Mohan et al 2010; Gamel et al 2005).

All samples show retrogradation of starch in the cooling phase (from 95 to 50°C). Retrogradation was much lower in sprouted chickpea flour, because the increased dextrinization by alpha amylase- limits starch reorganization as the suspension is cooled. The setback values were 479 and 314 BU for raw and

1
2
3 249 sprouted chickpea flour, respectively. This is an encouraging result in view of preparing chickpea enriched
4
5 250 bread, since low setback values are related to low starch retrogradation and thus, to improved shelf life.
6
7 251
8
9 252 **Dough enrichment with sprouted chickpea flour improves dough formation and leavening properties**
10
11 253 **compared to raw chickpea flour.** Wheat dough enrichment with sprouted chickpea determined important
12
13 254 changes in its dough-mixing behavior as measured by the Farinograph (Fig. 4A). Incorporation of raw
14
15 255 chickpea flours did not affect the optimal water absorption, that increased slightly when including sprouted
16
17 256 chickpea flour. A similar effect was observed by Fernandez and Berry (1989). Differences between raw and
18
19 257 sprouted chickpea enriched dough might be attributed to the formation of peptides and to the overall
20
21 258 proteins structural changes observed upon sprouting.
22
23
24 259 Dough development time is the time from the first addition of water to reaching the point of
25
26 260 greatest torque. During this phase of mixing, water hydrates the flour components and dough is developed.
27
28 261 High level of fiber usually results in longer development time (Peressini and Sensidoni 2009). Addition of
29
30 262 sprouted chickpea flour lowers development time much more than the non-sprouted one (5.4 vs 8.7 min).
31
32 263 These differences could relate to the effects of sprouting on thiol group accessibility. Increase in mixing
33
34 264 time of wheat dough with incorporation of either raw or sprouted chickpea (from 2 min of wheat flour to
35
36 265 8.7 min and 5.4 min for dough with raw and sprouted chickpea flours respectively) was in agreement with
37
38 266 previous studies (Baik and Han 2012; Mohammed et al 2012; Mohammed et al 2014; Sathe et al 1981).
39
40 267 The dough sample containing flour from sprouted chickpea exhibits a similar stability similar than
41
42 268 non-sprouted chickpea flour (7.5 vs 8.3 min). Despite the initial loss of consistency, the dough with
43
44 269 sprouted chickpea flour was not subjected to further sagging, maintaining torque values very similar to
45
46 270 unsprouted flour. The changes in dough characteristics upon addition of chickpea flour may be attributed
47
48 271 to dilution of gluten-forming proteins causing weakening of dough. Variation in hydration behavior of two
49
50 272 proteins may be another reason for these differences (Mohammed et al 2012; Mohammed et al 2014).
51
52
53
54
55 273 Dough leavening properties are reported in Fig. 4B and Fig. 4C, presenting dough height
56
57 274 development and gas production and retention during proofing, respectively. Interestingly, chickpea
58
59
60

enrichment increases dough development during proofing (from 39.2 mm in wheat-only dough to about 47.7 mm in and chickpea-enriched dough). Enrichment with sprouted chickpea gives a similar height as non-sprouted (47.7 vs 46.5 mm), but in a shorter time (103 vs 166 min), likely due to the sucrose formation promoted by sprouting (Table I).

The enrichment in sprouted chickpea flour allows the dough to reach the maximum development faster than in the case of non-sprouted flour (Fig. 4A). Addition of chickpea flour increases gas production from 1200 ml for wheat-only to 1538 ml (non-sprouted) and 1574 ml (sprouted). The amount of retained gas is higher in enriched dough compared to the control (1135 ml for wheat-only, 1380 ml (non-sprouted) and 1397 ml (sprouted)). The relation between gas production and retention, given as a percentage of gas retained in the dough, was slightly decreased by enrichment going from ~ 95 for wheat-only to 90 % (non-sprouted) and 89% (sprouted). This parameter is related to the dough ability to be stretched in thin membranes and - again - is associated to quality of the protein network. All together, these results suggest that partial germination of chickpea promotes a more rapid dough development during proofing, without worsening the gas production and retention of the dough.

CONCLUSIONS

This study proves that sprouted chickpeas can be used as an ingredient for production of enriched food. Enrichment with industrial-scale germinated chickpea flour appears to exert beneficial effects from a nutritional standpoint, whereas the structural modifications in macromolecules make sprouted chickpea flour a promising and interesting ingredient in formulating enriched products. Thus, sprouted chickpea flour offers a good opportunity for new products aiming at improving the nutritional significance of products targeted to low-income consumers and to users relying on vegetarian or vegan diets.

ACKNOWLEDGMENTS

The Authors thank Mr. Gaetano Cardone for technical assistance.

LITERATURE CITED

1
2
3 302 AACC International. Approved Methods of Analysis, 11th Ed. 2001. AACC International: St Paul, MN.
4
5 303 Abu-Ghannam, N., Gowen, A. 2011. Pulse-based food products. Pages 249-282 in: Pulse Foods. B. K.
6
7 304 Tiwari, A. Gowen and B. McKenna, eds. Elsevier Inc: Oxford, UK.
8
9 305 AOAC International. Official Methods of Analysis. 16th Ed. Method 991.43. 1995. Total, Insoluble
10
11 306 and Soluble Dietary Fiber in Food—Enzymatic-Gravimetric Method, MES-TRIS Buffer. AOAC International,
12
13 307 MD.
14
15 308 Arnon, R. 1970. Papain. Pages 226-244 in: Methods in Enzymology. G.E. Perlmann, ed. Academic
16
17 309 Press: New York and London.
18
19 310 Aslam, J., Mohajir, M.S., Khan, S.A, Khan, A.Q. 2008. HPLC analysis of water-soluble vitamins (B1,
20
21 311 B2, B3, B5, B6) in in vitro and ex vitro germinated chickpea (*Cicer arietinum* L.). African J. Biotech. 7: 2310-
22
23 312 2314.
24
25 313 Baik, B. K., Han, I. H. 2012. Cooking, roasting, and fermentation of chickpeas, lentils, peas, and
26
27 314 soybeans for fortification of leavened bread. Cereal Chem. 89:269-275.
28
29 315 Barbiroli, A., Bonomi, F., Casiraghi, M. C., Iametti, S., Pagani, M. A., Marti, A. 2013. Process
30
31 316 conditions affect starch structure and its interactions with proteins in rice pasta. Carbohydr. Polym. 92:1865-
32
33 317 1872.
34
35 318 Bellaio, S., Kappeler, S., Zamprogna Rosenfeld, E., Jacobs, M. 2013. Partially germinated ingredients
36
37 319 for naturally healthy and tasty products. Cereal Foods World 58:55-59.
38
39 320 Bellaio, S., Zamprogna Rosenfeld , E., Jacobs, M. 2011. Novel process based on partial germination
40
41 321 to enhance milling yield and nutritional properties of pulses. Cereal Foods World 56:A30.
42
43 322 Bellaio, S., Zamprogna Rosenfeld, E., Jacobs, M., Basu, S., Keppeler, S. 2012. Pargem, the technology
44
45 323 for a new family of healthy, safe, and convenient food ingredients based on partial germination. Cereal
46
47 324 Foods World 57:A3.
48
49 325 Bonomi, F., D'Egidio, M.G., Iametti, S., Marengo, M., Marti, A., Pagani, M.A. and Ragg, E.M. 2012.
50
51 326 Structure–quality relationship in commercial pasta: A molecular glimpse. Food Chem. 135:348-355.
52
53 327 Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of
54
55 328 protein utilizing the principle of protein–dye binding. Anal. Biochem. 72:248-254.
56
57
58
59
60

- 329 Cabrera-Chávez, F., de la Barca, A. M. C., Islas-Rubio, A. R., Marti, A., Marengo, M., Pagani, M. A.,
330 Bonomi, F., Iametti, S. 2012. Molecular rearrangements in extrusion processes for the production of
331 amaranth-enriched, gluten-free rice pasta. Food Sci. Technol.- LEB 47:421-426.
- 332 Chang, K. C., Harrold, R. L. 1988. Changes in selected biochemical components in vitro protein
333 digestibility and amino acids in two bean cultivars during germination. J. Food Sci. 53:783-787.
- 334 Chitra, U., Singh, U., Rao, P. V. 1996. Phytic acid, in vitro protein digestibility, dietary fiber, and
335 minerals of pulses as influenced by processing methods. Plant Foods Hum. Nutr. 49:307-316.
- 336 Collado, L. S., Corke, H. 1999. Accurate estimation of sweetpotato amylase activity by flour viscosity
337 analysis. Journal Agr. Food. Chem. 47:832-835.
- 338 Duenas, M., Sarmiento, T., Aguilera, Y., Benitez, V., Molla, E., Esteban, R. M., Martin-Cabrejas, M.A.
339 2016. Impact of cooking and germination on phenolic composition and dietary fibre fractions in dark beans
340 (*Phaseolus vulgaris* L.) and lentils (*Lens culinaris* L.). LWT-Food Sci. Techn. 66: 72-78.
- 341 Fernandez, M.L., Berry, J.W. 1989. Rheological properties of flour and sensory characteristics of
342 bread made from germinated chickpea. Int. J. Food Sci. Tech. 24:103-110.
- 343 Frohlich, P., Boux, G., Malcolmson, L. 2014. Effect of processing conditions during the partial
344 germination of whole yellow peas on the quality of spaghetti and extruded snacks. Cereal Foods World
345 59:A39.
- 346 Gamel, T.H., Linssen, J.P., Mesallem, A., S., Damir, A.A., Shekib, L.A..2005. Effect of seed treatments
347 on the chemical composition and properties of two amaranth species: starch and protein. J. Sci. Food Agr.
348 85:319-327.
- 349 Ghavidel, R. A., Prakash, J. 2007. The impact of germination and dehulling on nutrients,
350 antinutrients, in vitro iron and calcium bioavailability and in vitro starch and protein digestibility of some
351 legume seeds. Food Sci. Technol.- LEB 40:1292-1299.
- 352 Ghumman, A., Kaur, A., Singh, N. 2016. Impact of germination on flour, protein and starch
353 characteristics of lentil (*Lens culinari*) and horsegram (*Macrotyloma uniflorum* L.) lines. LWT-Food Sci. Tech.
354 65:137-144.

1
2
3 355 Gujral, H. S., Angurala, M., Sharma, P., Singh, J. 2011. Phenolic content and antioxidant activity of
4
5 356 germinated and cooked pulses. *Int. J. Food Prop.* 14: 1366-1374.
6
7 357 Han, J., Buchko, A. 2014. Development of a partial germination process for yellow peas and
8
9 358 resultant pea flours in white layer cakes. (Abstr.) *Cereal Foods World* 59:A42.
10
11 359 Iametti, S., Bonomi, F., Pagani, M. A., Zardi, M., Cecchini, C., & D'Egidio, M. G. 2006. Properties of
12
13 360 the protein and carbohydrate fractions in immature wheat kernels. *J. Agric. Food Chem.* 54:10239-10244.
14
15
16 361 ISO. Standard 14902. Animal feeding stuffs - Determination of trypsin inhibitor activity of soya
17
18 362 products. 2001.
19
20
21 363 Jain, A.K., Kumar, S., Panwar, J.D.S. 2009. Antinutritional factors and their detoxification in pulses- a
22
23 364 review. *Agric. Rev.* 30:64-70.
24
25 365 Khandelwal, S., Udipi, S. A., Ghugre, P. 2010. Polyphenols and tannins in Indian pulses: Effect of
26
27 366 soaking, germination and pressure cooking. *Food Res. Int.* 43:526-530.
28
29 367 Mahadevamma, S., Tharanathan, R. N. 2004. Processing of legumes: Resistant starch and dietary
30
31 368 fiber contents. *J. Food Qual.* 27:289-303.
32
33 369 Marti, A., Seetharaman, K., Pagani, M. A. 2013. Rheological approaches suitable for investigating
34
35 370 starch and protein properties related to cooking quality of durum wheat pasta. *J. Food Qual.* 36:133-138.
36
37 371 Martín-Cabrejas, M. A., Ariza, N., Esteban, R., Mollá, E., Waldron, K., & López-Andréu, F. J. 2003.
38
39 372 Effect of germination on the carbohydrate composition of the dietary fiber of peas (*Pisum sativum* L.). *J.*
40
41 373 *Agric. Food Chem.* 51:1254-1259.
42
43 374 Martín-Cabrejas, M. A., Díaz, M. F., Aguilera, Y., Benítez, V., Mollá, E., Esteban, R. M. 2008.
44
45 375 Influence of germination on the soluble carbohydrates and dietary fibre fractions in non-conventional
46
47 376 legumes. *Food Chem.* 107:1045-1052.
48
49
50 377 Mohammed, I., Ahmed, A. R., Senge, B. 2012. Dough rheology and bread quality of wheat–chickpea
51
52 378 flour blends. *Ind. Crop. Prod.* 36:196-202.
53
54 379 Mohammed, I., Ahmed, A. R., Senge, B. 2014. Effects of chickpea flour on wheat pasting properties
55
56 380 and bread making quality. *J Food Sci Technol.* 51:1902-1910.
57
58
59
60

- 381 Mohan, B.H., Malleshi, N.G., Koseki, T. 2010. Physicochemical characteristics and non-starch
382 polysaccharide contents of Indica and Japonica brown rice and their malts. *Food Sci. Technol.* 43:784-791.
- 383 Oberleas, D., Harland, B. 2007. Validation of a column liquid chromatographic method for phytate.
384 *J. AOAC Internat.* 90:1635 -1638.
- 385 Oomah, B. D., Patras, A., Rawson, A., Singh, N., Compos-Vega, R. 2011. Chemistry of pulses. Pages
386 9-55 in: *Pulse Foods*. B. K. Tiwari, A. Gowen and B. McKenna, eds. Elsevier Inc: Oxford, UK.
- 387 Peressini, D., Sensidoni, A. 2009. Effect of soluble dietary fibre addition on rheological and
388 breadmaking properties of wheat doughs. *J. Cereal Sci.* 49:190-201.
- 389 Sathe, S. K., Ponte, J. G., Jr., Rangnekar, P. D., and Salunkhe, D. K. 1981. Effects of addition of great
390 northern bean flour and protein concentrates on rheological properties of dough and baking quality of
391 bread. *Cereal Chem.* 58:97-100.
- 392 Savelkoul, F. H. M. G., Van der Poel, A. F. B., Tamminga, S. 1992. The presence and inactivation of
393 trypsin inhibitors, tannins, lectins and amylase inhibitors in legume seeds during germination. A review.
394 *Plant Foods Hum. Nutr.* 42:71-85.
- 395 Scarafoni, A., Consonni, A., Galbusera, V., Negri, A., Tedeschi, G., Rasmussen, P., Magni, C. and
396 Duranti, M. 2008. Identification and characterization of a Bowman-Birk inhibitor active towards trypsin but
397 not chymotrypsin in *Lupinus albus* seeds. *Phytochemistry* 69:1820-1825.
- 398 Vidal-Valverde, C., Frias, J., Sierra, I., Blazquez, I., Lambein, F., Kuo, Y. H. 2002. New functional
399 legume foods by germination: effect on the nutritive value of beans, lentils and peas. *Eur. Food Res. and*
400 *Technol.* 215:472-477.
- 401 Zamproga Rosenfeld, E., Bellaio, S., Mane, D., Jacobs, M. 2011. A new family of healthy, safe, and
402 convenient food products based on partial germination of pulses. *Cereal Foods World* 56:A71.
- 403 Zygmunt, L.C., Anderson, E., Behrens, B., Bowers, R., Bussey, M., Cohen, G., Colon, M., Deis, C.,
404 Given, P.S., Granade, A., Harms, C., Heroff, J.C., Hines, D., Hung, G.W., Hurst, W.J., Keller, J., Laroche, F.B.,
405 Luth, W., McKay, D., Mertle, T., Navarre, M., Rivera, R., Scopp, R., Scott, F., Sherman, R., Sloman, K.,
406 Sodano, C., Trick, K.D., Vandine, B.R., Webb N.G. 1982. High pressure liquid chromatographic determination
407 of mono and disaccharides in presweetened cereals: collaborative study. *J. Assoc. Anal. Chem.* 65:256-264.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

408 Figure legends

409 **Figure S1.** Sprouted chickpeas.

410 **Figure 1.** Effects of sprouting on trypsin inhibitory activity.

411 **Figure 2.** SDS-PAGE separation under non-reducing (top) and reducing (bottom) conditions of proteins
412 solubilized from sprouted and non-sprouted chickpeas in different media.

413 **Figure 3.** Effect of sprouting on pasting properties of chickpea flour.

414 **Figure 4.** Effect of sprouting on mixing properties (A), dough development (B) and gas production (C) of
415 chickpea-enriched dough.

416

Table I

Effect of sprouting on the chemical composition of chickpea flour

Data are expressed as g (100 g flour d.b.)⁻¹

	non-sprouted	sprouted
Starch	45.6 ± 1.3	42.1 ± 3.1
Protein	18.6 ± 0.3	20.2 ± 0.1
Lipid	7.3 ± 0.3	7.2 ± 0.6
Ash	3.2 ± 0.1	2.8 ± 0.0
Sugars	5.6 ± 0.1	5.6 ± 0.2
Glucose	0.1 ± 0.0	n.d
Fructose	0.1 ± 0.0	0.1 ± 0.0
Sucrose	3.2 ± 0.2	5.5 ± 0.2
Raffinose	2.2 ± 0.2	n.d.
Total Fiber	17.9 ± 1.7	18.8 ± 2.0
SolubleFiber	1.0 ± 0.1	1.6 ± 0.7
InsolubleFiber	16.9 ± 0.3	17.2 ± 0.1

TABLE II

Unbound micronutrient and phytic acid content in sprouted and non-sprouted chickpea flour.

Data are expressed as micrograms (g flour d.b.)⁻¹

analyte	non-sprouted	sprouted
Na	444.6 ± 10.0	442.8 ± 12.0
Mg	546.0± 16.0	455.0 ± 18.0
Ca	230.0 ± 10.0	263.0 ± 7.0
Cr	0.2±0.0	0.2 ± 0.0
Mn	1.6± 0.1	0.9± 0.0
Fe	3.9 ± 0.2	4.6 ± 0.1
Co	0.05 ± 0.01	0.08 ± 0.00
Ni	0.8 ± 0.1	1.0 ±0.0
Cu	0.4 ±0.0	1.4 ± 0.1
Zn	10.1 ± 0.3	9.9 ± 0.2
Se	0.1 ±0.0	0.2 ± 0.0
Mo	3.9 ± 0.2	4.2 ± 0.1
vit B ₁	0.4 ±0.1	0.8 ± 0.1
vit B ₂	0.4 ± 0.1	0.4 ± 0.0
vit B ₆	4.9 ± 0.2	5.3 ± 0.4
vit PP	0.7 ± 0.0	0.8± 0.0
phytic acid	18810 ± 107	17010 ± 279




Figure 1. Effects of sprouting on trypsin inhibitory activity.




Figure 2. SDS-PAGE separation under non-reducing (top) and reducing (bottom) conditions of proteins solubilized from sprouted and non-sprouted chickpeas in different media.






Figure 3. Effect of sprouting on pasting properties of chickpea flour.






Figure 4. Effect of sprouting on mixing properties (A), dough development (B) and gas production (C) of chickpea-enriched dough.



Figure S1. Sprouted chickpeas.

Peer Review